

中国弯螺属一新种记述（肺螺亚纲，柄眼目，扭轴蜗牛科）

周卫川<sup>1</sup> 张卫红<sup>2</sup> 陈德牛<sup>3</sup>

1. 福建出入境检验检疫局 福州 350001
2. 新疆大学生命科学与技术学院 乌鲁木齐 830046
3. 中国科学院动物研究所 北京 100101

摘 要 笔者在整理福建地区陆生贝类标本时，经比对鉴定发现 1 新种，即蛹形弯螺 *Sinoennea pupoidea* sp. nov.。隶属肺螺亚纲、柄眼目、扭轴蜗牛科、弯螺属。对新种形态特征、栖息环境作了记述，并对其近似种也进行了讨论。  
关键词 前鳃亚纲，柄眼目，扭轴蜗牛科，弯螺属，新种。  
中图分类号 Q959.212.5

蛹形弯螺，新种 *Sinoennea pupoidea* sp. nov. (图 2~ 5)

高0.57 mm, 壳口宽 0.57, 标本产自福建省龙岩龙洞 (25.30° N, 117.16° E)，2008 年 5 月 10 日。副

正模标本 壳高2.94 mm, 壳宽 1.68 mm, 壳口




图 1~ 5 蛹形弯螺，新种 *Sinoennea pupoidea* sp. nov.  
1. 贝壳侧面观 (lateral view of shell) 2 贝壳右侧面观 (right lateral of shell) 3 贝壳左侧面观 (left lateral of shell)  
4. 贝壳背侧面观 (dorsal lateral of shell) 5. 壳口放大 (aperture enlarge)

福建省自然科学基金项目 (2009J0104)、国家科技基础性工作专项 (2006FY111000- 02) 资助。  
收稿日期: 2008-07-22, 修订日期: 2009-07-16.

模标本 57 个, 壳高 2.80~3.00 mm, 壳宽 1.66~1.74 mm, 壳口高 0.55~0.63 mm, 壳口宽 0.54~0.63 mm。标本采自福建省永泰县天门山 (25.87°N, 118.96°E), 2008 年 4 月 26 日。福建省龙岩龙洞 (25.30°N, 117.16°E), 2008 年 5 月 10 日。福建省龙岩莲花山 (25.01°N, 117.05°E), 2008 年 5 月 11 日。福建省武夷山自然保护区 (27.75°N, 117.65°E), 2009 年 7 月 16 日。

模式标本采集地点如图 1 所示。

正模标本及一部分副模标本保存于福建出入境检验检疫局检验检疫技术中心标本室, 另一部分副模标本保存于中国科学院动物研究所标本馆。

**形态特征** 贝壳小型, 壳质厚, 结实, 有光泽, 呈蛹形, 有  $5\frac{1}{2}$  个螺层, 各螺层均匀增长, 膨大, 螺旋部高。壳面为乳白色或淡黄褐色, 壳顶 1~2 个螺层下壳面具有无数纵行排列的肋纹, 胚螺层光滑, 无肋纹。壳顶十分钝, 缝合线深。体螺层不膨大, 靠近壳口处逐渐变窄, 缩小。壳口狭小, 呈椭圆形,

口缘厚, 外翻, 扩大而延长, 呈喇叭状。壳轴垂直, 在壳口内唇上有 1 枚较大呈片状角板, 在右侧口缘上有 2 枚小的乳头状齿, 在轴缘处有 1 枚较大的乳头状齿。脐孔小, 呈孔隙状。

**栖息环境** 一般生活在阴暗潮湿多腐殖质石灰岩山区和丘陵地带, 常栖息于多地衣、苔藓的石灰岩石壁上或缝隙中, 落叶、腐败树枝下。新种在福建龙岩龙洞 (石灰岩地貌) 种群密度较高, 而在福建永泰天门山 (火山岩地貌) 种群密度较低。

**讨论** 新种与福州弯螺 *Sinoemea fuzouensis* (Zhou and Chen, 2006), 在外部形态上相似, 但较后者小, 新种有  $5\frac{1}{2}$  个螺层 (壳高 2.94 mm, 壳宽 1.68 mm, 壳口高 0.57 mm, 壳口宽 0.57), 在轴缘处有 1 枚较大的乳头状齿。但后者有  $7\frac{1}{2}$  个螺层 (壳高 5.17 mm, 壳宽 2.57 mm, 壳口高 1.86 mm, 壳口宽 1.48 mm), 在轴缘处无较大的乳头状齿。故而有所区别。




图 6 蛹形弯螺地理分布

Fig. 6. Geographical distribution of *Sinoemea pupidea* sp. nov.

## REFERENCES (参考文献)

Gredler, V. 1881. Zur Conchylienfauna von China III., Nachr. Bl. 118, pl. 6, f. 4.

Gredler, V. 1881. Zur Conchylienfauna von China VI., Nachr. Bl. 118, p. 223.

Heude, P. M. 1882. Notes sur les mollusques terrestres de la vallée du Fleuve Bleu., Mem. D'Hist. Nat. Emp. Chinois. 75, pl. 18, f. 23.

Luo, T-C and Chen, D-N 1998. Studies on *Sinoemea* Kobelt from China.

- Guizhou Sci.*, 16 (2): 90-93. [罗泰昌, 陈德牛, 1998. 中国弯螺属的研究. 贵州科学, 16 (2): 90~93]
- Luo, T. C., Zhou, W. C. and Chen, D. N. 2004. A new species of the *Sinoennea* Kobelt from China. (Pulmonata, Styommatophora, Streptaxidae). *Acta Zootaxonomica Sinica*, 29 (1): 77-79. [罗泰昌, 周卫川, 陈德牛, 2004. 中国弯螺属一新种 (肺螺亚纲, 柄眼目, 扭轴蜗牛科). 动物分类学报, 29 (1): 77~79]
- Moellendorff, O. V. 1881. Beiträge zur Molluskenfauna von Suedchina. Jahrb. 311.
- Moellendorff, O. V. 1882. Diagnoses specierum novarum chinae meridionalis. Jahrb. 183.
- Moellendorff, O. V. 1885. Materialien zur Fauna von China. Die Auriculaceen. Jahrb. 376.

- Moellendorff, O. V. 1887. Ueber die Enner Arten China's. Jahrb. 28.
- Panha, Somsak and Burch, J. B. 2005. An introduction to the microsnails of Thailand. *Malac. Review*, 127-129.
- Thiele, J. 1931. Handbuch der Systematischen Weichtierkunde. *Zool. Jahrb. Syst.*, 1: 149-151.
- Yen, T. C. 1939. Die Chinesischen land und Suesswasser Gastropoden des Natur Museums Senkenberg. *Abh. Senken. Nat. Ges.*, 444: 160-162, pl. 16, f. 36-43.
- Zhou, W. C., Chen, D. N. and Guo, Y. H. 2006. A new species of the *Sinoennea* Kobelt from Fujian, China (Pulmonata, Styommatophora, Streptaxidae). *Acta Zootaxonomica Sinica* 31 (2): 321-323. [周卫川, 陈德牛, 郭云海, 2006. 福建弯螺属一新种记述 (肺螺亚纲, 柄眼目, 扭轴蜗牛科). 动物分类学报, 31 (2): 321~323]

## A NEW SPECIES OF THE GENUS SINOENNEA FROM CHINA (PULMONATA, STYOMMATOPHORA, STREPTAXIDAE)

ZHOU Wei Chuan<sup>1</sup>, ZHANG Wei Hong<sup>2</sup>, CHEN De Niu<sup>3</sup>

1. Fujian Entry-Exit Inspection & Quarantine Bureau, Fuzhou 350001, China
2. College of Life-Science and Technology, Xinjiang University, Urumqi 830046, China
3. Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

**Abstract** A new species of the genus *Sinoennea* is described in the present paper. The materials were collected in Fujian Province, China in 2008 by the authors. Type specimens are deposited in the Fujian Entry-Exit Inspection & Quarantine Bureau, Fuzhou and the Institute of Zoology, Chinese Academy of Sciences, Beijing, China.

*Sinoennea pupoidea* sp. nov. (Figs 2-5)

Holotype, alt. 2.94 mm, diam. 1.68 mm, alt of aperture 0.57 mm, diam. of aperture 0.57 mm, collected from Longkoutong Town, Longyan City (25.30°N, 117.16°E), Fujian Province, China, 10 May 2008. Paratypes 57 specimens, alt. 2.80-3.00 mm, diam. 1.66-1.74 mm, alt of aperture 0.55-0.63 mm, diam. of aperture 0.54-0.63 mm, collected from Longkoutong Town, Longyan City (25.30°N, 117.16°E), 10 May 2008, Lianhuashan Town, Longyan City (25.01°N, 117.05°E), 11 May, 2008 and Tianmenshan Town, Wuyishan Nature Protection Areas (27.75°N, 117.65°E), 16 July 2009, Yongtai County (25.87°N, 118.96°E), 26 Apr. 2008, Fujian Province, China.

Geographical distribution of *Sinoennea pupoidea* sp. nov. is showed in Fig. 1.

**Key words** Pulmonata, Styommatophora, Streptaxidae, *Sinoennea*, new species.

Shell small, thick, solid, smooth, lustrous, pupa-shaped. Whorls  $5\frac{1}{2}$ , moderately convex with high spiral part. Shell milk white or light yellow in colour, and with some curve vertically rib-stria, but the embryonic and second without rib-stria, smooth. Apex blunt and short, ovate, with deep suture. Aperture small, elliptical-shaped. The peristome thick, expanded and reflexed, with 1 big flake-shaped angular lamellar at inner lip, with two nipple-formed teeth in right lip, with 1 big nipple-formed tooth in columellar lip. Umbilicus small, cavernous.

The new species fairly resembles *Sinoennea fuzouensis* (Zhou and Chen, 2006), but differs from the latter by the shells with  $5\frac{1}{2}$  whorls, (alt 2.94 mm, diam.

1.68 mm, alt of aperture 0.57 mm, diam. of aperture 0.57 mm) and with 1 big nipple-formed tooth in columellar lip. The latter shell with  $7\frac{1}{2}$  whorls, (alt 5.17 mm, diam. 2.57 mm, alt of aperture 1.86 mm, diam. of aperture 1.48 mm) and without 1 big nipple-formed tooth in columellar lip.